



HOLOCENE ECOLOGICAL TRAJECTORIES IN LAKE AND WETLAND SYSTEMS (AUVERGNE, FRANCE): A PALAEOENVIRONMENTAL CONTRIBUTION FOR A BETTER ASSESSMENT OF ECOSYSTEM AND LAND USE 'S VIABILITY IN MANAGEMENT STRATEGIES

Yannick Miras, Marlène Lavrieux, Marta Flórez

► To cite this version:

Yannick Miras, Marlène Lavrieux, Marta Flórez. HOLOCENE ECOLOGICAL TRAJECTORIES IN LAKE AND WETLAND SYSTEMS (AUVERGNE, FRANCE): A PALAEOENVIRONMENTAL CONTRIBUTION FOR A BETTER ASSESSMENT OF ECOSYSTEM AND LAND USE 'S VIABILITY IN MANAGEMENT STRATEGIES. *Annali di Botanica*, 2013, 3, pp.127-133. hal-00824420

HAL Id: hal-00824420

<https://hal.science/hal-00824420>

Submitted on 22 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HOLOCENE ECOLOGICAL TRAJECTORIES IN LAKE AND WETLAND SYSTEMS (AUVERGNE, FRANCE): A PALAEOENVIRONMENTAL CONTRIBUTION FOR A BETTER ASSESSMENT OF ECOSYSTEM AND LAND USE 'S VIABILITY IN MANAGEMENT STRATEGIES

MIRAS Y.^{1,2*}, LAVRIEUX M.^{3,4,5,6}, FLOREZ M.⁷

¹CNRS, GEOLAB, UMR 6042, Laboratoire de Géographie physique et environnementale,
4 rue Ledru, F-63057 Clermont-Ferrand, France

²Clermont Université, Université Blaise Pascal, GEOLAB, Maison des Sciences de l'Homme,
BP 10448, F-63000 Clermont-Ferrand, France

³Institut des Sciences de la Terre d'Orléans, Université d'Orléans, CNRS/INSU, BRGM,
ISTO, UMR 7327, 45071 Orléans, France.

⁴GéHCo, Géohydrosystèmes Continentaux, EA 6293, Faculté des Sciences et Techniques,
Université François Rabelais de Tours, Parc Grandmont, 37200 Tours, France

⁵Laboratoire des Sciences du Climat et de l'Environnement, CEA - Orme des Merisiers, 91191 Gif-sur-Yvette Cedex, France

⁶Chrono-Environnement, La Bouloie, UFR Sciences et Techniques, 16 route de Gray, 25030 Besançon Cedex, France

⁷Maison des Sciences de l'Homme de Clermont-Ferrand (USR 3550/CNRS), Université Blaise Pascal,
4 rue Ledru, 63057 Clermont-Ferrand cedex 1, France

*Corresponding author: Telephone: +33 473346822; e-mail: yannick.miras@univ-bpclermont.fr

(RECEIVED 04 MARCH; RECEIVED IN REVISED FORM 11 APRIL; ACCEPTED 12 APRIL)

ABSTRACT – Multi-proxy palaeoenvironmental studies are very important for the development of sustainable management strategies for ecosystems and modern landscapes. Analysing the relationship between societies, climate and environments through time, these studies contribute to define adequate policies and strategies for socio-environmental management, protection and legacy. Two complementary case studies– Aydat lake and Espinasse fen - from the south of the Chaîne des Puys (Auvergne, Massif Central, France) are presented. The analysis of these sedimentological records (both lacustrine and peat) follows a multi-proxy approach combining abiotic and biotic palaeoindicators (density, magnetic susceptibility, X-Ray Fluorescence spectrometry, Rock-Eval, pollen, non-pollen palynomorphs, molecular biomarkers). Aydat lake and Espinasse fen analyses were performed following a high spatio-temporal resolution. Results underline that long-term models of detrital input and eutrophication correspond to complex patterns with early and recurrent phases of human-induced ecological disturbances. They also evidence the existence of diversified long-term land use systems (deforestation, grazing, agriculture, hemp culture and retting) that provide fresh insights into the understanding of present-day mountain environments. This history between diversified human activities and hydrosystems responses must be taken into account for the construction of accurate retrospective and prospective model simulations of hydrosystem functioning.

KEYWORDS: HUMAN IMPACT, SOCIO-ENVIRONMENTAL VIABILITIES, EUTROPHICATION, DETRITAL INPUT, HYDROSYSTEMS, HOLOCENE, AUVERGNE

INTRODUCTION

Nowadays, a consensus exists within the global scientific community for underlining the importance of palaeoenvi-

ronmental studies in the development of strategies for the sustainable management of ecosystems and modern landscapes (Dearing et al., 2011). Palaeoenvironmental studies allow the understanding of the genesis and evolution

of ecosystems as well as the long-term shaping of landscape (e.g. temporal trajectories, thresholds). Through the analysis of the relationship between societies, climate and environments along time, they also contribute to define adequate policies and strategies for socio-environmental management, protection and legacy. Taking into account diachronic socio-environmental dynamics appears crucial in areas presenting a high sensitivity to global change such as the Mediterranean basin (e.g. Curras et al., 2012; Mercuri & Sadori, 2011) or European mountains (e.g. Ejarque et al., 2009; Galop et al., 2011). The mid-altitude mountainous configuration of the volcanic Chaîne des Puys (Auvergne, Massif Central, France), at the boundary of oceanic and continental climatic influences, makes this area particularly sensitive to climatic change. Besides, its vicinity with the regional capital city of Clermont-Ferrand explains a more remarkable anthropisation than in other parts of the Auvergne region. Finally, the socio-economic development of this region needs to be consistent with environmental legacy objectives such as the future registration of the

Chaîne des Puys at the UNESCO's World heritage list (<http://www.chainedespuys-failledelimage.com/>). Scientific stakes are thus important, particularly for recent ecological investigations at the Chaîne des Puys which aim at developing and testing new methods to design models that could guarantee the viability of both ecosystem quality (e.g. lake eutrophication) and land-use development in the area (e.g. touristic and agro-pastoral activities, residential areas) (Deffuant & Gilbert, 2011). Viability can be here defined as the persistence of multi-constrained system dynamics (Aubin, 1991), such as hydrosystems (e.g. lakes and wetlands). In this sense, diachronic palaeoecological studies are excellent tools since they allow (1) to describe a “pre-impact” state; (2) to draw the diachronic ecological processes; (3) to characterize forcing of disturbances and (4) to define a “post-impact” state and the mechanisms that drive resilience and human adaptability.

Two complementary case studies located at the south of the Chaîne des Puys –the Aydat lake (N 45°39.809' / E 2°59.106' / 837 m a.s.l.) and the Espinasse fen (N 45°38' / E 2°53' /




Figure 1. Location of Aydat lake and Espinasse fen (after Lavrieux et al., in press).

1160 m a.s.l.) - will be presented here to illustrate how palaeoecological data can be helpful in the proposed objectives and approach (Figure 1). They have been selected among the available palaeoenvironmental studies- including palynological analysis- which have been performed in this area because they have been carried out using a multi-proxy approach combining abiotic and biotic indicators and they propose the overall goal of analyzing hydrosystem's evolution along time with a particular interest on human impact (Lavrieux, 2011; Lavrieux et al., accepted; Lavrieux et al., in press; Miras et al., 2004). Indeed, previous studies were either mainly focused on vegetation history and tephrostratigraphy (Beaulieu & Goeury, 1987; Juvigné et al., 1988) or not carried out at a sufficient high temporal resolution (Michelin et al., 2001). In addition, recent investigations were not performed on long-term lake or fen records (Ballut et al., 2008; Ballut et al., 2012). Conversely, the 19 and 5-meter-long sedimentary records retrieved in the Aydat lake and the Espinasse fen respectively cover the last 6700 and 5000 yrs. Analyses were performed at a very high resolution and reliable age-depth models were constructed (Lavrieux et al., accepted; Lavrieux et al., in press; Miras et al., 2004). Besides, the close vicinity of these two sites, which are located in the same drainage basin, allows a comparison of environmental changes at a micro-local scale. Micro-local studies are indeed the most suitable approach to analyse human-environments interactions (Dearing et al., 2006), particularly in upland environments (Ejarque et al., 2010). The Aydat lake and the Espinasse fen are currently two anthropogenic hotspots in the Auvergne region. Both sites are highly frequented by tourists and account for agro-pastoral activities (Figure 2), and as a result of that, they are deeply impacted by human activities. For instance, the eutrophication level of Lake Aydat can be so high that summer swimming is sometimes forbidden. In consequence,

local authorities, organizations of environmental management (e.g. Parc Naturel Régional des Volcans d'Auvergne) and the scientific community have begun to pay attention to the restoration of the ecosystem services, combined with a sustainable socio-economic development. This paper aims to demonstrate that palaeoecological investigations have to be taken into account as they furnish essential data concerning the long-term anthropogenic forcing on hydrosystem evolution. This paper is focused in: (1) model of sedimentation and (2) eutrophication phases.

MATERIALS AND METHODS

Multi-proxy approach is based on the integration of abiotic (density, magnetic susceptibility, X-Ray Fluorescence spectrometry, Rock-Eval) and biotic indicators (pollen, non-pollen palynomorphs, and molecular biomarkers). Employed methods follow standard procedures (see complete methodology in Lavrieux et al., accepted; Lavrieux et al., in press; Miras et al., 2004). Age-depth models are based on AMS radiocarbon dates: 17 for the Aydat lake and 5 for the Espinasse fen. In the Aydat lake, the top of the sequence has been dated using ^{137}Cs , ^{241}Am , ^{210}Pb and ^7Be measurements and the correlation with historical flood events (Lavrieux et al., accepted).



Figure 2. Current situation of Aydat lake (a. photo M. Lefeuvre) and Espinasse (b. photo Y. Miras).

RESULTS

Sedimentation and erosive mass wasting

Sediment analysis of Aydat lake's reveals two units on the basis on sedimentological and bulk geochemical indicators (Lavrieux et al., accepted). Multi-proxy palaeoenvironmental analyses were performed on the lower (1076-1974 cm depth) and the upper unit (0-829 cm depth). The intermediate unit (829-1076 cm depth) corresponds to a mass wasting deposit (Lavrieux et al., accepted). Between 6730 and 3200 cal BP, 6 phases of relatively higher background detrital input have been underlined, the first one starting at ca 6000 cal BP with a major flood event reported at ca 3800 cal BP. In the Espinasse fen, detrital phases are also observed between ca 4850 and 3900 cal BP (Miras et al., 2004). This first Aydat unit ends with an erosive mass wasting deposit dated to ca 1770±60 cal BP. From 1770 cal BP to the present day, 6 phases of higher background detrital input are distinguished in the Aydat lake's sediments and dated at ca 1510-1210, 1130-1000, 950-670, 620-500, 430-150 and 90-50 cal BP (Lavrieux et al., accepted).

Eutrophication phases

Different trophic states have been evidenced in the Aydat lake on the basis of pollen and non-pollen palynomorph data and sediment geochemical data. Earlier phases of trophic variation and/or nutrient enrichment occurred between 6000-5750 and 4900-4600 cal BP. Eutrophication phases have also been highlighted during the lapses 3900-3500 and 1470-888 cal BP, reaching a maximum level between 1200 and 1130 cal BP (Lavrieux et al., accepted) and 515 and 200 cal BP (work in progress). The recent period— between 150 cal BP/present day- also corresponds to a high nutrient level.

DISCUSSION

Both the Aydat lake and the Espinasse fen sequences reveal long-term processes of detrital input and changes in eutrophic conditions (Figures 3 and 4). Pollen and non-pollen palynomorph data indicate that this variability in lake status correspond to early and recurrent phases of human-induced ecological disturbances. Together with a climate driver— lower solar activity could favour enhanced soil erosion on catchments (Debret et al., 2010), the anthropogenic forcing is also evidenced for the phases at ca 6000-5700 and 4800-4600 cal BP (Lavrieux et al., accepted). Moreover, ecological changes dated to 3800 cal BP are not correlated with a

climatic event but with a more intense landscape management (3900-3500 cal BP). This period corresponds to the major flood event in the Aydat sequence and to an erosive mass wasting in Espinasse fen (Lavrieux et al., accepted; Miras et al., 2004). Pollen and charcoal data suggest a clear fire-induced woodland opening associated with the extension of grasslands and heliophilous taxa. The significant rise of anthropogenic pollen indicators and dung-related fungi attest the expansion of local grazing activity which provokes the first nutrient enrichment of the lake. If climatic conditions and human impact worked together until the Mid-Holocene, sedimentation and eutrophication were principally controlled by anthropogenic forcing during the Late Holocene, especially since 1500 cal BP. During historical times, background sedimentation increases and over-enrichment phases of the Aydat lake are always synchronous with more intensive and persistent human pressure on environment. All this indicates that lake and fen systems in the Chaîne des Puys have intensively and continuously been impacted by human activities since the Neolithic and the Early Bronze age and these mountains must not been considered as pristine and marginal areas. These ancient anthropogenic impacts also underline the high vulnerability of these ecosystems. Such a long-term history of human activities and hydrosystem interactions must thus be taken into account for the construction of accurate retrospective and prospective simulations models of hydrosystem functioning.

Multi-proxy palaeoecological investigations performed in the Southern part of the Chaîne des Puys also evidence complex and diversified land use systems along time. This contrasts with a common view of European mountain areas which tends to focus on ecological disturbances uniquely caused by grazing activity. Even if this practice had an important impact on upland hydrosystems since millennia, this is only one of the multiple activities which were developed in mountain areas (Ejarque et al., 2011). In the South of the Chaîne des Puys, pollen data evidence that agriculture reaches at the altitudinal of the Aydat lake in determinate phases. This is the case for the period from the 12th to the 18th centuries (work in progress). Moreover, pollen and the cannabinoil molecular tracer indicate that hemp culture and retting activity were also very important, particularly between the 13th and the 19th centuries (Lavrieux et al., in press). However, a hemp culture is also suggested during Gallo-roman times. Hemp retting caused a degradation of the water quality of the Aydat lake during the Modern period, as suggested by recurrent blooms of *Anabaena* akinetes (work in progress). This concept of a diversified exploitation of natural resources in mountain environments undoubtedly provides fresh insights into the understanding of mountain present-day environments.




Figure 3. Pollen and non-pollen palynomorphs percentage diagram of selected taxa from Aydat Lake (after Lavrieux et al., in press).

CONCLUSIONS

The overall objective of this paper was demonstrating that lessons can be learned from multi-proxy palaeoenvironmental investigations. Results obtained can be of benefit for the long-term sustainability and the management of ecosystems and services. The design of mitigation strategies to both current and future socio-environmental stresses requires a good understanding of the long-term response of ecosystems to cumulative changes caused by climate variations and human activities. This integrated and transdisciplinary approach also addresses the adaptability of past societies to environmental (e.g. climatic) changes and furnishes valuable tools to manage consequences of global change. This research is still in progress and a better quantitative and spatial assessment of past human impact is required. For this, two research strategies can here be quoted. Firstly, new palaeoecological methods need to be tested to obtain valuable and innovative biotic indicators, mainly within the

non-pollen palynomorphs assemblages, for exploring new questions such as the nature of grazing, eg. type of animal, grazing system, or different land-use systems. Secondly, this integrated palaeo- and ecological approach should be – if possible! – supplemented with archaeological, historical and ethnographical data-sets in order to properly evaluate, characterize and spatialize human occupation and land use. This work is actually in progress in the Chaîne des Puys.

ACKNOWLEDGEMENTS

The Aydat lake study was supported by the Erode project funded by the INSU/CNRS. We would like to thank all the members involved in this project (ISTO, GéHCo, LSCE, IMBE, Edytem) as well as thus involved in the multi-proxy study of the Espinasse fen (ISTO, Chrono-environnement,




Figure 4. Pollen percentage diagram of selected taxa from Espinasse fen (after Miras et al., 2004).

IMBE). Many thanks also to Dr A. Ejarque (Northern Arizona University, Flagstaff) for her precious comments and corrections of the English of the text. Finally, we would like to thank Dr M. Lefevre who is the author of the Aydat Lake photo.

REFERENCES

- Aubin J.-P., 1991. Viability theory, Birkhäuser.
- Ballut C., Michelin Y., Miras Y., 2012. Landscape human shaping and spatial mobility of the agropastoral practices in the Chaîne des Puys during the historical times (French Massif central). *Quaternary International* 251, 97-106.
- Ballut C., Prat B., Lopez Saez J.-A., Gaby G., Cabanis M., 2008. Evolution environnementale d'une zone humide et de son bassin versant depuis la fin de l'âge du Fer: le maar de

- Montchâtre (Massif central, France). *Quaternaire* 19(1), 87-97.
- de Beaulieu J.-L., Goeury C., 1987. Zonation automatique appliquée à l'analyse pollinique : exemple de la narse d'Ampoix (Puy de Dôme). *Bulletin de l'Association française pour l'Etude du Quaternaire* 29, 49-61.
- Currás A., Zamora B., Reed J.M., García-Soto E., Ferrero S., Armengol X., Mezquita-Joanes F., Marqués M.A., Riera S., Julià R., 2012. Climate change and human impact in central Spain during Roman times: high-resolution multi-proxy analysis of a tufa lake record (Somolinos, 1280 m asl). *Catena* 89, 31-53.
- Dearing J., Dotterweich M., Foster T., 2011. Editorial: integrative paleoscience for sustainable management. *PAGES news* 19(2), 43.
- Dearing J.A., Battarbee R.W., Dikau R., Larocque I., Oldfield F., 2006. Human–environment interactions: learning from the past. *Regional Environmental Change* 6, 115-23.
- Debret M., Chapron E., Desmet M., Rolland-Revel M., Magand O., Trentesaux A., Bout-Roumazielle V., Nomade J., Arnaud F., 2010. Northwestern Alps Holocene paleohydrology recorded by flooding activity in Lake Le Bourget, France. *Quaternary Science Reviews* 29, 2185-2200.
- Deffuant G., Gilbert N., 2011. *Viability and Resilience of Complex Systems: Concepts, Methods and Case Studies from Ecology and Society*. Springer.
- Ejarque A., Julià R., Riera S., Palet J.M., Orengo H.A., Miras Y., Gascón C., 2009. Tracing history of highland human management in the Eastern Pre-Pyrenees (Spain): an interdisciplinary palaeoenvironmental approach. *The Holocene* 19(8), 1241-1255.
- Ejarque A., Miras Y., Riera Mora S., 2011. Modern pollen and non-pollen palynomorph analogues of vegetation and highland grazing activities obtained from surface and dung datasets from the eastern Pyrenees. *Review of Palaeobotany and Palynology* 167, 123-139.
- Ejarque A., Miras Y., Riera S., Palet J.M., Orengo H. 2010. Testing microregional variability in the Holocene shaping of high mountain cultural landscape: a palaeoenvironmental case-study in the Eastern Pyrenees. *Journal of Archaeological Science* 37, 1468-1479.
- Galop D., Houet T., Mazier F., Leroux G., Rius D., 2011. Grazing activities and biodiversity in the Pyrenees: new insight on high altitude ecosystems in the framework of a Human-Environment Observatory. *PAGES news* 19(2), 53-55.
- Juvigné E., Milcamps V., Delibrias G., Evin J., 1988. Ages de traits polliniques et chronozonation du Tardiglaciaire et de l'Holocène dans le Massif central (France). *Medelingen rijks geologisch dienst* 41(4), 33-50.
- Lavrieux M., 2011. *Biomarqueurs moléculaires d'occupation des sols, du sol au sédiment : exemple du bassin-versant et du lac d'Aydat (Puy-de-Dôme)*. Thesis, University François-Rabelais, Tours, France, 246 p.
- Lavrieux M., Jacob J., Disnar J.R., Bréheret J.G., Le Milbeau C., Miras Y., Andrieu-Ponel V., 2013. Sedimentary cannabinol tracks the history of hemp retting. *Geology*, in press.
- Lavrieux M., Disnar J.R., Chapron E., Bréheret J.G., Jacob J., Miras Y., Reyss J.L., Andrieu-Ponel V., Arnaud F., accepted. 6,700-year sedimentary record of climatic and anthropogenic signals in Lake Aydat (French Massif Central). *The Holocene*.
- Michelin Y., Vergne V., Cougoul C., Cournut S., 2001. Variations des teneurs en éléments minéraux dans un bas marais holocène: la Vézolle (Chaîne des Puys), première recherche des manifestations anthropiques. *Quaternaire* 1(2), 31-41.
- Miras Y., Laggoun-Défarge F., Guenet P., Richard H., 2004. Multi-disciplinary approach to changes in agro-pastoral activities since the Subboreal in the surroundings of the "narse d'Espinasse" (Puy de Dôme, French Massif Central). *Vegetation History and Archaeobotany* 13, 91-103.
- Mercuri A.M., Sadori L., 2013. Mediterranean culture and climatic change: past patterns and future trends. In: Goffredo S., Dubinsky Z. (Eds.), *The Mediterranean Sea: Its History and Present Challenges*. Springer, Dordrecht.